

Euclidean vs. non-Euclidean Gamma-Ray Bursts

M. Tavani

IFCTR-CNR, via Bassini 15, I-20133 Milano (Italy)

Columbia Astrophysics Laboratory, Columbia University, New York, 10027 NY (USA)

Abstract

We classify gamma-ray bursts (GRBs) according to their observed durations and physical properties of their spectra. We find that long/hard bursts (of duration $T_{90} > 2.5$ s, and typical photon energy $E_p \gtrsim 0.8$ MeV corresponding to BATSE's energy fluence hardness $H_{32}^e > 3$) show the strongest deviation from the three-dimensional Euclidean brightness distribution. The majority of GRBs, i.e., short bursts ($T_{90} < 2.5$ s) and long/soft bursts (with $T_{90} > 2.5$ s, and $H_{32}^e < 3$) show little, if any, deviations from the Euclidean distribution. These results contradict the prediction of extragalactic GRB models that the most distant bursts should be the most affected by cosmological energy redshift and time-dilation (long/soft GRBs). The strongly non-Euclidean GRB subclass has very hard spectra of typical photon energy above ~ 1 MeV, i.e., outside the ideal energy range for optimal detection by BATSE. Selection effects due to BATSE inefficient triggering and/or intrinsic burst properties may cause the strong deviation from the Euclidean distribution for the hardest class of long GRBs. Our results imply strong luminosity and spectral evolution of GRB sources placed at extragalactic distances. Whether there co-exist multiple GRB populations with different (Galactic or extragalactic) origins remains an open question.

Submitted to the *Astrophysical Journal Letters*: November 24, 1997

Revised: 27 January 1998; accepted: 13 February 1998

Subject Headings: Gamma-rays: bursts, theory

1. Introduction

The brightness distribution of gamma-ray bursts (GRBs) depends on both the intrinsic properties of the emission and the spatial distribution of sources. When analyzed globally, GRBs detected by BATSE show a clear deviation from the three-dimensional Euclidean brightness distribution (Meegan et al. 1992, 1995). However, recent analyses of special GRB subclasses showed that the deviation from the Euclidean distribution does not apply to all bursts. On the contrary, ‘soft’ bursts with typical long durations appear to have an Euclidean brightness distribution (Pizzichini 1995, Kouveliotou 1996, Belli 1997, Pendleton et al. 1997). A commonly used spectral parameter is the *burst-averaged* hardness ratio H_{32} , i.e., the ratio of overall fluences in the BATSE energy channels no. 2 and 3 corresponding to the 100-300 keV and 50-100 keV bands, respectively. The photon fluence-averaged ratio H_{32}^p , and the energy fluence-averaged ratio H_{32}^e are commonly used. A first approach was to consider different GRB subclasses ranked according to their H_{32}^e (e.g., Pizzichini 1995). The analysis of Kouveliotou et al. (1996) found that GRBs with photon-fluence-averaged $H_{32}^p > 4$ and $H_{32}^p < 2$ are distributed quite differently, with the soft bursts showing an Euclidean distribution. Belli (1997) used a different phenomenological criterion to distinguish GRB classes in the hardness/duration diagram (H_{32}^e vs. T_{90} , with T_{90} the duration of 90% energy deposition in the 50-300 keV band) divided by the relation $H_{32}^e = 2 T_{90}^{1/2}$. The long-duration subclass with $H_{32}^e > 4.5$ shows the most pronounced deviation from the Euclidean distribution.

Pendleton et al. (1997, hereafter P97) used a very different approach reaching qualitatively similar conclusions. P97 select GRB *individual pulses* according to the existence or lack of a high energy component above 300 keV (BATSE’s channel no. 4). Two classes of GRBs with without a high energy component (HE and NHE) were identified. Very often, complex GRBs with multiple pulses show both kinds of HE and NHE pulses.

The interesting conclusion of this analysis is that NHE pulses, both individual or as part of more complex GRBs, appear to have Euclidean distributions (P97).

All of these analyses were carried out on a purely phenomenological level, with no direct correspondence between the selecting criterion and a physical model of emission. The main motivation of our work is to provide a physical model for the GRB observable quantities (T_{90}, H_{32}) to better understand the GRB brightness distribution. We use the shock synchrotron model (SSM) of GRB emission that successfully explains the great majority of GRB burst-averaged spectra (Tavani 1996a,b, 1997; hereafter T96a,b, T97). The results of our GRB analysis are obviously independent of the SSM: only the interpretation of the adopted GRB selection depends on the theoretical model. We studied GRBs of the Fourth BATSE (4B) Catalog (Paciesas et al. 1997). Three effects should be considered: (1) the intrinsic spectral evolution within bursts (typically hard-to-soft) that tends to concentrate bursts diagonally from top left to bottom right of the $H_{32}^e - T_{90}$ diagram; (2) the effect due to cosmological redshift, or any other cooling mechanism simultaneously dilating the time profile and softening the spectrum (T96a,b); (3) selection effects due to BATSE sensitivity. If the majority of GRB sources are at extragalactic distances, an overall softening of dim and time-dilated bursts is expected as these bursts should be the most distant for no spectral cosmological evolution. The population of soft and long duration GRBs is expected to be inhomogeneously distributed if BATSE’s sampling radius exceeds the spatial distance of peak burst formation rate. As we show below, this is the opposite of what we find.

2. Physical interpretation of T_{90} and H_{32}

The double-peaked distribution of GRB durations is well known (Fishman & Meegan 1995), and probably reflects different manifestations of simple (short) and complex (long) burst emission. Pulse analysis of complex bursts shows a structure of pulses with average

durations of $\sim 200 - 400$ ms (Norris et al. 1996). The subclass of short bursts with $T_{90} < 2.5$ s can be modelled as relatively simple bursts whose substructure consists of a single (or a few) pulse(s) of average properties. Longer bursts ($T_{90} > 2.5$ s) are either apparently dilated versions of simple short bursts, or complex superpositions of many pulses of average properties. The double-peaked T_{90} distribution can then be related to the relative complexity of the pulse structure. No apparent correlation exists between overall burst durations and peak photon intensities. We define here ‘short’ and ‘long’ bursts divided by the critical duration $T_{90} = 2.5$ s corresponding to the local minimum of the T_{90} distribution.

There is instead a definite correlation between T_{90} and average GRB hardness, most likely as a consequence of the hard-to-soft spectral evolution within bursts (e.g., Ford et al., 1995). Fig. 1 shows the 4B hardness/duration diagram (H_{32}^e vs. T_{90}) with a clear indication that short bursts are generally harder than long ones. Spectral softening for individual pulses and for long bursts has been reported in many cases (Ford et al 1995), and it constitutes a fundamental property of cooling particle distributions (e.g., Tavani, 1998).

We apply the synchrotron shock model (T96a,b) for the interpretation of the hardness ratio H_{32}^e . The SSM is based on a hydrodynamic and radiative model of relativistic MHD outflows interacting with shocks that reconvert part of their energy in impulsive acceleration. The SSM can in principle be applied to relativistic mass/electromagnetic outflow models at extragalactic and Galactic distances. Average hardness ratios H_{32}^e can be reliably converted into average peak energies, E_p , of the νF_ν spectrum (T97). Fig. 3 of T97 shows the SSM calculated H_{32}^e (and H_{32}^p) as a function of E_p for different assumptions about the index δ of the high-energy component of the post-acceleration particle energy distribution function ($N(\gamma) \propto \gamma^{-\delta}$). The SSM calculated H_{32}^e ’s are in good agreement with detailed broad-band average spectra of GRBs (T96a,b;T97). The agreement between theory and observations extends also to GRBs detected above 1 MeV (T96a,b). a factor of ~ 2 with the SSM calculated value an apparent depletion of soft photons We can therefore

physically interpret the average hardness ratio by converting its value to a typical photon energy E_p at which most of the burst energy is dissipated. We choose $H_{32}^e = 3$ as the critical value distinguishing GRBs with typical photon energies below or above 0.8 MeV. We call them ‘soft’ and ‘hard’ bursts, respectively. We also carried out calculations for different choices of the critical $H_{32,c}^e$, in the range $2.5 \lesssim H_{32,c}^e \lesssim 4$ that would correspond to different peak energies (~ 400 keV and ~ 2 MeV, respectively; see Fig. 3 of T97). Our main conclusions do not depend on the choice of $H_{32,c}^e$ in the range given above. Our preferred choice ($HR_{32,c}^e = 3$) is physically justified by selecting GRBs whose typical E_p is ~ 4 times the average E_p detected by BATSE (~ 200 keV, Band et al. 1993), and outside the optimal energy range for BATSE detection (50-300 keV). The energy spectrum of GRB 910503 (Schaefer et al. 1996, T96a) clearly shows the spectral properties of GRBs corresponding to our critical value for $H_{32,c}^e$. The range for ideal BATSE detection is $1 \lesssim H_{32}^e \lesssim 2.5$.

Fig. 1 clearly shows that short GRBs have average E_p well above 1 MeV, and are efficiently detected by a combination of short risetimes (below 1 s) and spectral evolution in the BATSE’s optimal detection range. We notice the relative small number of short/soft GRBs, contrary to the large number of long/soft bursts. This feature is not influenced by selection effects, and reflects the GRB hard-to-soft spectral evolution from simple GRBs to multiple-pulsed events. Selection effects may play a role in determining the trigger efficiency for long/hard events populating the upper right corner of the hardness/duration diagram. In our analysis, we consider only bursts with well defined spectral properties, including only events with relative uncertainty on H_{32}^e less than 50%. We also note that only $\sim 3\%$ of the events in Fig. 1 show a deviation of their measured H_{32}^e by more than 3σ larger than the ideal maximum allowed SSM value ($H_{32}^e \simeq 5$, see T97). These events are most likely characterized by soft photon depletion/absorption in the 50-300 keV band that may influence the observed H_{32}^e in addition to instrumental effects.

3. Analysis of the 4B catalog

Equipped with a physical interpretation of the $H_{32}^e - T_{90}$ diagram, we analyzed the brightness distribution of 4B GRBs according to their different hardness ratios and durations. We distinguish four classes:

(S/H) GRBs with $T_{90} < 2.5$ sec, and $H_{32}^e > 3$ (short/hard bursts);

(S/S) GRBs with $T_{90} < 2.5$ sec, and $H_{32}^e < 3$ (short/soft bursts);

(L/H) GRBs with $T_{90} > 2.5$ sec, and $H_{32}^e > 3$ (long/hard bursts);

(L/S) GRBs with $T_{90} > 2.5$ sec, and $H_{32}^e < 3$ (long/soft bursts).

Fig. 2 shows the results, where we plot the cumulative burst number with peak photon intensity above a given value P_{64} as a function of P_{64} (64 ms trigger timescale). Both classes L/S and S/H show brightness distributions in good agreement with the $-3/2$ Euclidean slope. The deviations from the Euclidean distribution are minor, with a discrepancy below a factor of 2 between real and inferred values from the high- P_{64} 's at $P_{64} = 1.3 \text{ ph. cm}^{-2} \text{ s}^{-1}$. The curvature near $P_{64} \lesssim 1 \text{ ph. cm}^{-2} \text{ s}^{-1}$ is clearly due to a selection effect. Both L/S and S/H distributions might also be marginally consistent with a high-flux slope of $-3/2$ followed by an intermediate-flux slope near -1 for $P_{64} \lesssim 8 \text{ ph. cm}^{-2} \text{ s}^{-1}$. On the contrary, the long/hard (L/H) bursts show a strong deviation from the Euclidean distribution at $P_{64} \simeq 20 \text{ ph. cm}^{-2} \text{ s}^{-1}$. The fourth class of short/soft events contains very few (66) and faint events and its brightness distribution is not relevant here¹. These results are confirmed also for brightness distributions obtained for 256 and 1024 ms trigger timescales.

All four duration/hardness GRB classes show isotropic sky distributions. Table 1 gives the values obtained for the dipole and quadrupole moments relative to the Galactic center

¹ If we summed together all the soft and hard events with $T_{90} < 2.5$ s we would obtain results very similar to those for the S/H class.

and plane. We also obtained the values of the V/V_{max} parameter for GRBs with tabulated ratios C_p/C_{min} (with C_p and C_{min} the peak and threshold countrates, respectively for the 64 ms trigger timescale in the BATSE’s energy range). Table 1 reports the average values of V/V_{max} for the three duration/hardness GRB classes. We note that a spectral bias in the determination of C_{min} is to be expected (see below), and that tabulated values of C_{min} can be lower limits to the actual minimum countrates for detection of bursts with duration/spectral properties outside BATSE’s optimal range.

Our results indicate that only long/hard GRBs shows a strong deviation from the Euclidean brightness distribution. These events dominate the bright end of the GRB population (one out of two GRBs are L/H bursts in the range $6 \text{ ph. cm}^{-2} \text{ s}^{-1} \lesssim P_{64} \lesssim 100 \text{ ph. cm}^{-2} \text{ s}^{-1}$) and yet they show the strongest deviation from the Euclidean slope.

4. Discussion

We briefly discuss here some points relevant to the interpretation of our results, postponing a more thorough discussion to a later publication. Our results confirm previous analyses of the GRB brightness distribution, and contradict the predictions of simple cosmological models with no strong spectral and luminosity evolution. The total sample of GRBs detected by BATSE shows a clear deviation at low peak intensities from the Euclidean brightness distribution. This property, together with an average V/V_{max} value near 0.37 (see Table 1) was interpreted as a signature of a globally inhomogeneous distribution in space (e.g., Paczynski 1991, Briggs 1996). However, we find that this conclusion can be applied only to the special subclass of long/hard bursts. The fact that *only long/hard GRBs are strongly non-Euclidean being at the same time very intense* is puzzling for extragalactic models. The soft and long GRBs that would naturally exhibit

an inhomogeneous brightness distribution in simple cosmological models show little, if any, deviations from being Euclidean. We emphasize that our conclusion is deduced by the *combination* of average spectral and time dilation properties of GRBs. Furthermore, the peak intensity distributions of S/H, L/S and L/H bursts are similar to each other, making unnatural that L/H bursts are in the foreground of more distant (and possibly softer) bursts. We are left with the conclusion that strong luminosity and spectral cosmological evolution is necessary to reconcile our results with extragalactic models of GRBs. L/H bursts placed in remote regions where a relative decrease of their spatial density makes them strongly non-Euclidean would have to be on the average very bright (by a factor of several) and with harder spectra (E_p larger by a factor of ~ 10) than the majority of closer bursts. Whether this can be accomplished by GRB formation mechanisms in star forming galaxies (SFGs) (as suggested by recent HST observations of GRB 970508, Pian et al. 1998) remains to be seen. The redshift dependence of SFG density (Madau et al. 1996) is strongly peaked near $z \simeq 1$ and then decreases for larger redshifts. L/H bursts might originate at redshifts $z \gtrsim 1$ under the condition that more distant GRB sources produce bursts more luminous and harder than those at smaller redshifts. The V/V_{max} distribution of L/H bursts interpreted at face value with no selection bias, indicate that these bursts tend to reside in relatively close regions. The May 8th 1997 burst detected by BSAX (Piro et al. 1997) has an equivalent hardness $H_{32}^e = 3.6 \pm 0.9$ and $T_{90} = 8$ s (Amati et al. 1997). If GRB 970508 belongs to the L/H subclass of GRBs, is in the strongly non-Euclidean part of the brightness distribution. Our results imply that, if the optical transient at cosmological distance is associated with the burst, GRB 970508 might be located at a redshift larger than $z = 0.83$ (indicated by optical spectroscopy, Metzger et al. 1997).

Alternative interpretations of our results should also be considered. One possibility that reconciles our results with those of Pendleton et al. (1997) is that BATSE selection effects play a major role in shaping the brightness distribution of duration/hardness

subclasses of GRBs. The ability by BATSE to trigger on L/H events might crucially depend on their peak intensity (see also Lloyd & Petrosian 1997). As clearly indicated by the P97 results, even single pulses within complex GRBs share the same characteristics of our burst-averaged S/H, L/S and L/H subclasses. This fact demonstrates that short and long/soft bursts (pulses) can be detected with good efficiency given BATSE’s trigger conditions. However, detection of relatively long and hard bursts (or L/H pulses usually occurring at the beginning of complex GRBs, see P97) require special conditions. It is conceivable that these trigger conditions are favorably met for quite bright GRBs above $P_{64} \gtrsim 10 \text{ ph. cm}^{-2} \text{ s}^{-1}$. Detecting a burst (pulse) which is long/hard (or with substantial flux above 300 keV) may be difficult for smaller values of P_{64} . This might explain the sharp deviation from the Euclidean slope of the L/H burst brightness distribution. Since BATSE detects short/hard bursts with no apparent trigger inefficiency for low peak intensities (see Fig. 2), we conclude that long duration bursts (possibly with slow risetimes) with typical E_p ’s outside BATSE’s optimal detection range are the most affected by the trigger criteria. The results of Lloyd & Petrosian (1997) indicating a corrected E_p distribution with a large undetected population of GRBs with $E_p \gtrsim 1 \text{ MeV}$ agree with this interpretation. BATSE’s trigger conditions [flux enhancement over background nominally by 5.5 standard deviations within a given timescale (64, 256 and 1024 ms) and for the energy range 50-300 keV] strongly favors detection of short bursts as well as long bursts with typical emission in the optimal energy range (our L/S bursts). Slow risetime-bursts with average $E_p \gtrsim 1 \text{ MeV}$ and no strong hard-to-soft spectral evolution are disfavored for BATSE’s detection. *Both* burst-averaged and individual-pulse properties of GRB brightness distributions share the same Euclidean (non-Euclidean) character of their soft (long/hard) population. This fact finds a natural interpretation in BATSE’s selection effects.

We also note that BATSE spectral selection effects can strongly influence the actual values of C_{min} . Taken at face values, the average V/V_{max} ’s obtained for our duration/spectral

GRB classes (Table 1) indicate inhomogeneous distributions by more than three standard deviations for all except the S/S bursts. However, BATSE’s trigger conditions can couple the distance and spectral dependence of countrates in a non-trivial way, breaking the $1/r^2$ dependence of C_p deduced in previous applications of the V/V_{max} method to GRBs (e.g., Schmidt, Higdon & Hueter, 1988). The nominal values of C_{min} tabulated in the 50-300 keV range can underestimate (by a factor of a few) the actual values of C_{min} necessary to detect bursts well outside BATSE’s optimal spectral and risetime ranges. The long/hard and short/hard GRB populations are the most affected by this bias.

Disregarding a Galactic origin for GRBs might have been premature. The combination of sky isotropy, inhomogeneity and the average value of $V/V_{max} < 1/2$ applied to the whole sample of GRBs was interpreted as being inconsistent with standard Galactic models (e.g., Paczynski 1991). However, our results clearly indicate that the strong deviation from a homogeneous distribution (if that is the correct interpretation of non-Euclidean GRBs) applies only to long/hard bursts, and certainly not to the majority of GRBs. A Galactic origin of a subclass or the majority of GRBs is not excluded by our results.

The author thanks an anonymous referee for helpful comments.

References

- Amati L., and the BSAX Team, 1997, personal communication.
- Band, D., et al., 1993, ApJ, 413, 281.
- Belli, B.M., 1997, ApJ, 479, L31.
- Briggs, M., et al., 1996, ApJ, 459, 40.
- Fishman, G.J. & Meegan, C.A., 1995, ARA&A, 33, 415.
- Ford, L.A. et al., 1995, ApJ, 439, 307.
- Kouveliotou C., et al., 1996, AIP Conf. Proc. no. 384, p. 42.
- Lloyd, N.M. & Petrosian, V., 1997, in the Proceedings of the 4th Huntsville Gamma Ray Burst Symposium, eds. C.A. Meegan, P. Cushman (New York: AIP), in press (astro-ph/9711192).
- Madau, P. et al., 1996, MNRAS, 283, 1388.
- Meegan C.A., et al., 1992, Nature, 355, 143.
- Meegan, C.A., et al., 1995, 3B catalog, ApJS, 106, 65.
- Metzger, M.L., et al., 1997, Nature, 387, 878.
- Norris, J.P. et al., 1996, ApJ, 459, 393.
- Paciesas, W.S., et al., 1997, The Fourth BATSE Gamma-Ray Burst Catalog,
<http://www.batse.msfc.nasa.gov/data/grb/4bcatalog>.
- Pendleton, G. et al., 1997, ApJ, in press (P97).
- Pian E. et al., 1998, ApJ, 492, L103.

Piro L., et al., 1997, A&A, submitted.

Pizzichini, G., 1995, in the Proc. of the XXIV ICRC Conference, p. 81.

Schaefer B. et al., 1996, in AIP Conf. Proc. Series no. 384, p. 274.

Schmidt, M., Higdon, J.C. & Hueter, G., 1988, ApJ, 329, L85.

Tavani, M., 1996a, Phys. Rev. Letters, 76, 3478 (T96a).

Tavani, M., 1996b, ApJ, 466, 768 (T96b).

Tavani, M., 1997, ApJ, 479, 135 (T97).

Tavani, M., 1998, in the Proceedings of the 4th Huntsville Gamma Ray Burst Symposium,
eds. C.A. Meegan, P. Cushman (New York: AIP), in press.

Table 1: Properties of duration/hardness GRB subclasses

GRB duration/hardness class	$\langle \cos(\theta) \rangle$	$\langle \sin^2(b) - 1/3 \rangle$	$\langle V/V_{max} \rangle$
long/soft (520)	-0.002 ± 0.043	-0.007 ± 0.013	0.434 ± 0.019 (221)
long/hard (338)	-0.020 ± 0.054	$+0.008 \pm 0.016$	0.289 ± 0.022 (168)
short/hard (254)	-0.062 ± 0.063	-0.019 ± 0.018	0.354 ± 0.023 (149)
short/soft (66)	$+0.003 \pm 0.123$	$+0.042 \pm 0.036$	0.458 ± 0.044 (43)
total (1178)	-0.019 ± 0.016	-0.007 ± 0.013	0.373 ± 0.012 (581)

The table gives the dipole, quadrupole moments with respect to the Galactic center and plane, and the average value of V/V_{max} . We follow the definitions of Briggs et al. (1996). The angle θ is the angle between the burst position and the Galactic center. The angle b is the Galactic latitude.




Fig. 1.— GRB energy fluence-averaged hardness ratio (H_{32}^e) vs. duration (T_{90}) distribution from the 4th BATSE catalog (4B). Only GRBs with fractional error on H_{32}^e less than 50% are plotted. Error bars are 1σ . Long/hard (L/H) GRBs are marked by solid filled circles: they show the strongest deviation from the Euclidean brightness distribution (see Fig. 2).




Fig. 2.— Brightness distribution (cumulative number of bursts N above a certain peak photon intensity level P_{64} as a function of P_{64} (64 ms trigger timescale) for different subclasses of GRBs. *Solid histogram*: total distribution; *short-dashed histogram*: long/soft bursts; *long-dashed histogram*: long/hard bursts; *dotted-dashed histogram*: short/hard bursts.